



University of
New Haven

University of New Haven
Digital Commons @ New Haven

Mathematics Faculty Publications

Mathematics

1-2013

Contact Hypersurfaces of a Bochner-Kaehler Manifold

Amalendu Ghosh

Krishnagar Government College, India

Ramesh Sharma

University of New Haven, rsharma@newhaven.edu

Follow this and additional works at: <http://digitalcommons.newhaven.edu/mathematics-facpubs>



Part of the [Mathematics Commons](#)

Publisher Citation

Ghosh, A. & Sharma, R. (2013). Contact hypersurfaces of a Bochner-Kaehler manifold. *Results in Mathematics*, 64, 155-163. doi: 10.1007/s00025-013-0305-y

Comments

This is the authors' accepted manuscript. The final publication is available at Springer via <http://dx.doi.org/10.1007/s00025-013-0305-y>.

Contact Hypersurfaces Of A Bochner-Kaehler Manifold

Amalendu Ghosh¹ and Ramesh Sharma²

¹*Department Of Mathematics, Krishnagar Government College, Krishnanagar 741101, West Bengal, India.*

E-mail: aghosh_70@yahoo.com

²*Department Of Mathematics, University Of New Haven, West Haven CT 06516, USA.*

E-mail: rsharma@newhaven.edu

Abstract: We have studied contact metric hypersurfaces of a Bochner-Kaehler manifold and obtained the following two results: (1) A contact metric constant mean curvature (*CMC*) hypersurface of a Bochner-Kaehler manifold is a (k, μ) -contact manifold, and (2) If M is a compact contact metric *CMC* hypersurface of a Bochner-Kaehler manifold with a conformal vector field V that is neither tangential nor normal anywhere, then it is totally umbilical and Sasakian, and under certain conditions on V , is isometric to a unit sphere.

Keywords: Bochner-Kaehler manifold, Contact metric hypersurface, Constant mean curvature, Conformal vector field.

MS Classification: 53B25, 53C55, 53 C15.

1 Introduction

Bochner curvature tensor was introduced in 1948 by S. Bochner [4] as a Kaehlerian analogue of the Weyl conformal tensor. It was shown by S.M. Webster [17] that the fourth order Chern-Moser curvature tensor of *CR*-manifolds coincides with the Bochner tensor. A Kaehler manifold with vanishing Bochner curvature tensor is known as Bochner-Kaehler manifold. Bochner-Kaehler surface is nothing but a self-dual Kaehler surface in Penrose's twistor theory. Some topological obstructions to

Bochner-Kaehler metrics were studied by Chen in [6]. Just as a real space-form is conformally flat, a complex space-form is Bochner flat, i.e. Bochner-Kaehler (the converse does not need to hold). The product of two complex space-forms of constant holomorphic sectional curvatures c and $-c$ is non-Einstein Bochner-Kaehler. Though Bochner-Kaehler manifolds have been studied by quite a few geometers, nevertheless have received considerably less attention, compared to Kaehler metrics with vanishing scalar curvature and Kaehler-Einstein metrics. For details we refer to Bryant [5]. It is well known that a hypersurface M of a Kaehler manifold \bar{M} admits an almost contact metric structure induced from the Hermitian structure of \bar{M} . Okumura [13] studied and classified such hypersurfaces, mainly when the ambient space is a complex space-form. Generalizing the following result of Sharma [15] “The contact metric hypersurface of a complex space-form is a (k, μ) -contact manifold”, we prove the following main result of this paper.

Theorem 1 *A contact metric constant mean curvature hypersurface of a Bochner-Kaehler manifold is a (κ, μ) -contact manifold.*

Finally, we consider the case when the ambient space admits a conformal vector field and provide the following extrinsic characterization of a Sasakian manifold.

Theorem 2 *Let M be a compact contact metric constant mean curvature hypersurface of a Bochner-Kaehler manifold \bar{M} admitting a conformal vector field V which is neither tangential nor normal anywhere on M . Then M is Sasakian and totally umbilical in \bar{M} , and the component U of V , tangential to M is conformal on M . Further, (i) if U is non-Killing and $\dim.M > 3$, then M is isometric to the unit sphere S^{2n+1} , and (ii) if V is closed, then for any dimension, M is isometric to S^{2n+1} .*

2 Contact Metric Hypersurfaces Of A Kaehler Manifold

A $(2n + 1)$ -dimensional smooth manifold M is said to be a contact manifold if it carries a global 1-form η such that $\eta \wedge (d\eta)^n \neq 0$ everywhere on M . Given a contact 1-form η , there exists a unique vector field ξ such that $(d\eta)(\xi, X) = 0$ and $\eta(\xi) = 1$. Polarizing $d\eta$ on the contact subbundle

D ($\eta = 0$), one obtains a Riemannian metric g and a (1,1)-tensor field φ such that

$$(d\eta)(X, Y) = g(X, \varphi Y), \eta(X) = g(X, \xi), \varphi^2 = -I + \eta \otimes \xi \quad (1)$$

g is called an associated metric of η and (φ, η, ξ, g) a contact metric structure. The operators $h = \frac{1}{2}\mathcal{L}_\xi\varphi$ and $l = R(\cdot, \xi)\xi$ are self-adjoint and satisfy: $h\xi = 0$ and $h\varphi = -\varphi h$. Furthermore, $h, h\varphi$ are trace-free. Following formulas hold on a contact metric manifold.

$$\nabla_X \xi = -\varphi X - \varphi hX \quad (2)$$

$$l - \varphi l \varphi = -2(h^2 + \varphi^2) \quad (3)$$

If the associated CR-structure on M is integrable, then M is called a contact strongly pseudo-convex integrable CR manifold. This CR integrability condition was shown by Tanno [16] to be equivalent to

$$(\nabla_X \varphi)Y = g(X + hX, Y)\xi - \eta(Y)(X + hX) \quad (4)$$

and holds on a 3-dimensional contact metric manifold. A contact metric manifold (M, g) is said to be K -contact if ξ is Killing (equivalently, $h = 0$), and Sasakian if the almost Kaehler structure on the cone $M \times \mathbb{R}$ with metric $dr^2 + r^2g$ is Kaehler. Sasakian manifolds are K -contact and 3-dimensional K -contact manifolds are Sasakian. For details we refer to Blair [1]. In [2] Blair, Koufogiorgos and Papantoniou introduced a class of contact metric manifolds $M^{2n+1}(\eta, \xi, g, \varphi)$ satisfying the nullity condition:

$$R(X, Y)\xi = k(\eta(Y)X - \eta(X)Y) + \mu(\eta(Y)hX - \eta(X)hY) \quad (5)$$

for real constants k and μ . Such manifolds are known as (k, μ) -contact manifolds, and satisfy: $k \leq 1$, equality holding when M is Sasakian.

Let M be an isometrically embedded orientable hypersurface of a Kaehler manifold \bar{M} of real dimension $2n + 2$ and with complex structure tensor $J : J^2 = -I$ and the Hermitian metric g . The induced metric on M will also be denoted by g . If N denotes the unit normal vector field to M , we set

$$JN = \xi \quad (6)$$

$$JX = \varphi X - \eta(X)N, \quad (7)$$

where φ and η denote a $(1, 1)$ -tensor field and a 1-form respectively, and X an arbitrary vector field tangent to M . The Gauss and Weingarten formulas are

$$\bar{\nabla}_X Y = \nabla_X Y + g(AX, Y)N, \quad \bar{\nabla}_X N = -AX$$

where X, Y denote arbitrary vector fields tangent to M , ∇ and $\bar{\nabla}$ the Riemannian connections of M and \bar{M} respectively, and A the Weingarten operator. Differentiating (1) along an arbitrary vector field X tangent to M , using the Weingarten formula, and comparing tangential parts gives

$$\nabla_X \xi = -\varphi AX. \quad (8)$$

One can easily verify using (6) and (7) that (φ, ξ, η, g) defines the almost contact metric structure. We now assume that the almost contact metric structure induced on M is a contact metric structure. Using the formula (2) in (8) yields

$$A\xi = (Tr.A - 2n)\xi. \quad (9)$$

$$AX = X + hX + (Tr.A - 2n - 1)\eta(X)\xi. \quad (10)$$

which were derived in [15]. Next, differentiating (7) along M , and using (10) gives equation (4). Hence M is contact strongly pseudo-convex integrable CR manifold. We denote the Ricci tensor of M , of types $(0, 2)$ and $(1, 1)$ by S and Q respectively, and the scalar curvature by r of M . Corresponding objects of \bar{M} are denoted by the same letters with overbars. Recall the Gauss equation

$$\begin{aligned} g(\bar{R}(X, Y)Z, W) &= g(R(X, Y)Z, W) \\ &+ g(AX, Z)g(AY, W) - g(AY, Z)g(AX, W). \end{aligned} \quad (11)$$

and contract it as

$$\begin{aligned} \bar{S}(Y, Z) - g(\bar{R}(N, Y)Z, N) &= S(Y, Z) \\ &+ g(AX, AZ) - (Tr.A)g(AY, Z). \end{aligned} \quad (12)$$

For a Bochner-Kaehler manifold \bar{M} , the Bochner curvature tensor B (see [19]) vanishes, i.e. for arbitrary vector fields $\bar{X}, \bar{Y}, \bar{Z}, \bar{W}$ on \bar{M} , we have

$$\begin{aligned}
0 &= g(B(\bar{X}, \bar{Y})\bar{Z}, \bar{W}) = g(\bar{R}(\bar{X}, \bar{Y})\bar{Z}, \bar{W}) - \frac{1}{2n+6}[g(\bar{Y}, \bar{Z})g(\bar{Q}\bar{X}, \bar{W}) \\
&\quad - g(\bar{Q}\bar{X}, \bar{Z})g(\bar{Y}, \bar{W}) + g(\bar{J}\bar{Y}, \bar{Z})g(\bar{Q}\bar{J}\bar{X}, \bar{W}) - g(\bar{Q}\bar{J}\bar{X}, \bar{Z})g(\bar{J}\bar{Y}, \bar{W}) \\
&\quad + g(\bar{Q}\bar{Y}, \bar{Z})g(\bar{X}, \bar{W}) - g(\bar{X}, \bar{Z})g(\bar{Q}\bar{Y}, \bar{W}) + g(\bar{Q}\bar{J}\bar{Y}, \bar{Z})g(\bar{J}\bar{X}, \bar{W}) \\
&\quad - g(\bar{J}\bar{X}, \bar{Z})g(\bar{Q}\bar{J}\bar{Y}, \bar{W}) - 2g(\bar{J}\bar{X}, \bar{Q}\bar{Y})g(\bar{J}\bar{Z}, \bar{W}) \\
&\quad - 2g(\bar{J}\bar{X}, \bar{Y})g(\bar{Q}\bar{J}\bar{Z}, \bar{W}) + \frac{\bar{r}}{(2n+4)(2n+6)}[g(\bar{Y}, \bar{Z})g(\bar{X}, \bar{W}) \\
&\quad - g(\bar{X}, \bar{Z})g(\bar{Y}, \bar{W}) + g(\bar{J}\bar{Y}, \bar{Z})g(\bar{J}\bar{X}, \bar{W}) \\
&\quad - g(\bar{J}\bar{X}, \bar{Z})g(\bar{J}\bar{Y}, \bar{W}) - 2g(\bar{J}\bar{X}, \bar{Y})g(\bar{J}\bar{Z}, \bar{W})] \tag{13}
\end{aligned}$$

3 Proofs Of The Results

Lemma 1 *For a contact metric hypersurface of a Kaehler manifold,*

$$(a) \bar{S}(\varphi Y, Z) + \bar{S}(Y, \varphi Z) = \eta(Y)g(\bar{Q}N, Z) + \eta(Z)g(\bar{Q}N, Y)$$

$$(b) g(\xi, \bar{Q}N) = 0.$$

Proof: Since \bar{M} is Kaehler, we have $\bar{S}(JY, Z) + \bar{S}(Y, JZ) = 0$. The use of (7) in this gives (a). Substituting ξ for Y and Z in (a) yields (b).

Lemma 2 *If f is a smooth function on a contact metric manifold M such that $df = (\xi f)\eta$ (d denoting exterior derivation), then f is constant on M .*

Proof: Taking the exterior derivative of the differential condition mentioned in the hypothesis gives $d(\xi f) \wedge \eta + (\xi f)d\eta = 0$. Taking its wedge product with η we find $(\xi f)(d\eta) \wedge \eta = 0$. As $(d\eta) \wedge \eta$ is nowhere vanishing on M (otherwise the definition of contact structure would break down), we conclude that $\xi f = 0$ on M . Consequently, $df = 0$ on M , and hence f is constant on M , completing the proof.

Lemma 3 *For a contact metric hypersurface M of a Bochner-Kaehler manifold, the following conditions are equivalent:*

- (a) *For any vector field X tangent to M , $g(\bar{Q}N, X) = 0$*
- (b) *ξ is an eigenvector of the Ricci operator Q at each point of M*
- (c) *The mean curvature of M is constant.*

Proof : Using equations (12), (13), and part (b) of Lemma 1 gives

$$\begin{aligned}
(2n+5)\bar{S}(Y, Z) &= [\bar{S}(N, N) - \frac{\bar{r}}{2n+4}]g(Y, Z) - \frac{3\bar{r}}{2n+4}\eta(Y)\eta(Z) \\
&+ 3g(\bar{Q}Z, \xi)\eta(Y) + 3g(\bar{Q}Y, \xi)\eta(Z) + (2n+6)[S(Y, Z) \\
&+ g(AY, AZ) - (Tr.A)g(AY, Z)]
\end{aligned}$$

Now we replace Y by φY in the above equation to get one equation, and replace Z by φZ to get another equation. Adding these two equations, and using part (a) of Lemma 1 we obtain

$$\begin{aligned}
(2n+5)\{\eta(Y)g(\bar{Q}N, Z) + \eta(Z)g(\bar{Q}N, Y)\} &= 3g(\bar{Q}\varphi Z, \xi)\eta(Y) \\
+ 3g(\bar{Q}\varphi Y, \xi)\eta(Z) + (2n+6)[S(\varphi Y, Z) + S(Y, \varphi Z) + g(A\varphi Y, AZ) \\
+ g(AY, A\varphi Z) - (Tr.A)g(A\varphi Y, Z) - (Tr.A)g(AY, \varphi Z)]. & \quad (14)
\end{aligned}$$

Substituting ξ for Z , using (9) and part (b) of Lemma 1 yields

$$(n+1)\varphi\bar{Q}\xi = (n+3)\varphi Q\xi. \quad (15)$$

We also note

$$g(\bar{Q}N, X) = -g(J\bar{Q}\xi, X) = -g(\varphi\bar{Q}\xi, X), \quad (16)$$

The equations (15) and (16) show that (a) is equivalent to (b). Contracting the Codazzi equation: $\bar{R}(X, Y)N = (\nabla_Y A)X - (\nabla_X A)Y$ at X provides $\bar{S}(Y, N) = Y(Tr.A) - (div A)Y$. Equation (10) transforms the preceding equation into

$$\bar{S}(Y, N) = Y(Tr.A) - (\xi Tr.A)\eta(Y) - (div h)Y \quad (17)$$

Using equation (4) and the formula $(div.h)\xi = 0$ (easy to verify) for a contact metric shows

$$(div h\varphi)\varphi Y = -(div h)Y \quad (18)$$

Let us assume (b), i.e. $Q\xi = (Tr.l)\xi$. Applying the formula: $(div h\varphi)Y = S(Y, \xi) - 2n\eta(Y)$ (see [3]), we have $(div h\varphi)\varphi Y = S(\varphi Y, \xi) = 0$. Hence equation (18) shows that $div h = 0$. As $Q\xi = (Tr.l)\xi$ is equivalent to (a) [proven earlier], appealing to equation (17) we obtain $d(Tr.A) = (\xi Tr.A)\eta$. Application of Lemma 2 shows that $Tr.A$ is constant on M , proving (b) \Rightarrow (c). For the converse, assume (c), i.e. $Tr.A$ constant. Then, we go back to equations (17) and (18) and use the formula

$(\operatorname{div}h\varphi)Y = S(Y, \xi) - 2n\eta(Y)$ once again, getting $\bar{S}(X, N) = S(\varphi X, \xi) = -g(\varphi Q\xi, X)$. Using this in (16) we find $\varphi\bar{Q}\xi = \varphi Q\xi$. Finally, using this in (15) we conclude that $\varphi Q\xi = 0$ which implies (b), and complete the proof.

Lemma 4 *For a contact metric hypersurface M of a Bochner-Kaehler manifold \bar{M} ,*

$$(a)(Q\varphi - \varphi Q) - (\eta \circ Q\varphi) \otimes \xi + \eta \otimes \varphi Q\xi = 2(\operatorname{Tr}.A - 2)h\varphi.$$

$$(b)l\varphi - \varphi l = 2(\operatorname{Tr}.A - 2n)h\varphi.$$

Proof Replacing Y, Z by $\varphi Y, \varphi Z$ respectively, in (14) and then using (10) we get (a). Using the formula:

$$Q\varphi - \varphi Q = l\varphi - \varphi l + 4(n-1)h\varphi + (\eta \circ Q\varphi) \otimes \xi - \eta \otimes \varphi Q\xi = 0,$$

for a contact pseudo-convex integrable CR-manifold (see [10]), and using it in (a) we obtain (b).

Proof Of Theorem 1. First, we use equation (13) to obtain

$$\begin{aligned} g(\bar{R}(X, Y)\xi, W) &= \frac{1}{2n+6}[\eta(Y)g(\bar{Q}X, W) - g(\bar{Q}X, \xi)g(Y, W) \\ &\quad - g(\bar{Q}X, N)g(JY, W) + g(\bar{Q}Y, \xi)g(X, W) - \eta(X)g(\bar{Q}Y, W) \\ &\quad + g(\bar{Q}Y, N)g(JY, W) - 2g(JX, Y)g(\bar{Q}\xi, W)] \\ &\quad - \frac{\bar{r}}{(2n+6)(2n+4)}[\eta(Y)g(X, W) - \eta(X)g(Y, W)] \quad (19) \end{aligned}$$

for arbitrary vector fields X, Y, W tangent to M . By Lemma 3, the constant mean curvature hypothesis is equivalent to $g(\bar{Q}N, X) = 0$, i.e. $\bar{Q}N = fN$ for some function f on M . We also have $\bar{Q}\xi = J\bar{Q}N = f\xi$. Hence equation (19) reduces to

$$\begin{aligned} (2n+6)g(\bar{R}(X, Y)\xi, W) &= \eta(Y)g(\bar{Q}X, W) - \eta(X)g(\bar{Q}Y, W) \\ &\quad + \sigma[\eta(Y)g(X, W) - \eta(X)g(Y, W)] \quad (20) \end{aligned}$$

where $\sigma = f - \frac{\bar{r}}{2n+4}$. This shows that $g(\bar{R}(X, Y)\xi, W) = 0$, for any vector fields X, Y tangent to M and orthogonal to ξ . Next, substituting ξ for Z in (11), and using equations (9) and (20) we obtain $R(X, Y)\xi = 0$,

for any vector fields X, Y tangent to M and orthogonal to ξ . Hence by the result of Koufogiorgos-Stamatiou ([11]) we conclude that M is a (κ, μ) -space provided the dimension of M is ≥ 5 .

It remains to consider the 3-dimensional case for which we know that

$$\begin{aligned} R(X, Y)Z &= g(QY, Z)X - g(QX, Z)Y + g(Y, Z)QX \\ &\quad - g(X, Z)QY - \frac{r}{2}\{g(Y, Z)X - g(X, Z)Y\} \end{aligned} \quad (21)$$

Making use of the formula (3) and the formula $h^2 = (k - 1)\varphi^2$ (for any 3-dimensional contact metric manifold [9], where k is a function $= \frac{Tr.l}{2}$) in part (b) of Lemma 4, we obtain

$$lY = -k\varphi^2Y + (Tr.A - 2)hY \quad (22)$$

Differentiating this along an arbitrary vector field X , using (2) and then contracting the resulting equation at X with respect to a local orthonormal frame e_i , we find

$$\begin{aligned} &g((\nabla_Y Q)\xi, \xi) - g((\nabla_\xi Q)Y, \xi) - g(R(Y, \varphi e_i + \varphi h e_i)\xi, e_i) \\ &- g(R(Y, \xi)(\varphi e_i + \varphi h e_i), e_i) = -(\varphi^2 Y)\kappa + (Tr.A - 2)(div h)Y \end{aligned} \quad (23)$$

Using $Q\xi = (Tr.l)\xi$ and (2) we have $g((\nabla_Y Q)\xi, \xi) = 2(Yk)$, $g((\nabla_\xi Q)Y, \xi) = 2(\xi k)\eta(Y)$. We also had found during the proof of Lemma 3 that $div.h = 0$. Moreover, using (21) and $Q\xi = (Tr.l)\xi$ we compute

$$g(R(Y, \varphi e_i + \varphi h e_i)\xi, e_i) = -\eta(Y)Tr.(Q\varphi h)$$

and

$$g(R(Y, \xi)(\varphi e_i + \varphi h e_i), e_i) = 0$$

Utilizing all these findings in (23), we obtain

$$Yk - (\xi k)\eta(Y) + \eta(Y)Tr.(Q\varphi h) = 0$$

Taking $Y = \xi$ it is easy to see that $Tr.Q\varphi h = 0$. Hence $Yk = (\xi k)\eta(Y)$, i.e. $dk = (\xi k)\eta$. Applying Lemma 2, we conclude that k is constant. Thus, the hypothesis : $Q\xi = (Tr.l)\xi$ and (21) imply $R(X, Y)\xi = 0$, for any vector field X, Y orthogonal to ξ . Replacing X by $X - \eta(X)\xi$ and Y by $Y - \eta(Y)\xi$ (as these vector fields are orthogonal to ξ) we obtain

$$R(X, Y)\xi = \eta(Y)lX - \eta(X)lY$$

The use of (22) in the foregoing equation shows that M^3 is a (κ, μ) space with $\mu = (Tr.A - 2)$. This completes the proof.

Proof Of Theorem 2. As V is conformal on \bar{M} ,

$$\mathcal{L}_V g = 2\rho g \quad (24)$$

We decompose the conformal vector field V along M orthogonally as

$$V = U + \alpha N \quad (25)$$

where U is the tangential part of V and α a smooth function on M . In view of the Lemma 3, the constant mean curvature hypothesis is equivalent to $\bar{S}(X, N) = 0$ for arbitrary vector field X tangent to M . Following the procedure given on pages 101-104 of Yano [18], we have

$$2n \int_M \bar{S}(U, N) dM = \int_M \alpha \sum_{i \neq j}^{2n+1} (k_i - k_j)^2 dM \quad (26)$$

where dM is the volume element of M , and k_i are the principal curvatures of M . As the left hand side of the above equation is zero, and α is nowhere zero on M (otherwise V would become tangent to M somewhere, contradicting our hypothesis), we conclude that $k_i = k_j$, i.e. M is totally umbilical. Hence, using (10) provides $A = I$, and $h = 0$, i.e. M is Sasakian. The conformal Killing equation (24), together with the Gauss and Weingarten formulas show that $\mathcal{L}_U g = 2(\rho + \alpha)g$, i.e. U is conformal on M . If U is homothetic, then U reduces to Killing, since M is compact. Hence, if U is not Killing, and $dim > 3$, then by the following theorem of Okumura [14] "A complete Sasakian manifold of dimension > 3 and admitting a non-Killing conformal vector field is isometric to a unit sphere", M is isometric to S^{2n+1} which proves (i). For (ii), we know from the following result of Goldberg [8] "A closed conformal vector field on a non-flat Kaehler manifold is homothetic and holomorphic" that V is homothetic. Hence $\nabla_X V = \rho X$ (ρ constant). Using the decomposition (25), and bearing in mind that X is arbitrary tangent vector on M , we immediately obtain $U = -D\alpha$ (D is the gradient operator of (M, g)) and $\nabla_X U = (\alpha + \rho)X$. We note that α cannot be constant on M , otherwise U would vanish on M turning V normal to M and thus contradicting our hypothesis. Thus obtain

$$\nabla_X D(\alpha + \rho) = -(\alpha + \rho)X \quad (27)$$

Hence, by Obata's theorem [12] "A complete Riemannian manifold of dimension > 2 is isometric to a sphere of radius $\frac{1}{c}$ if and only if it admits a non-trivial solution f of the differential equation $\nabla\nabla f = -c^2fg$ ", we conclude that M is isometric to S^{2n+1} , completing the proof.

Remark. As indicated in [7], the Kaehler cone manifold $(M \times R^+, d(r^2\eta))$ with metric $dr \otimes dr + r^2g$ over a Sasakian manifold (M, η, g) admits a conformal vector field $ar\partial_r - b\xi$ (for a, b real constants) which is nowhere tangent and nowhere normal to M and therefore serves as an example of the conformal vector field satisfying the hypothesis of Theorem 2.

4 Acknowledgment:

R.S. was supported by the University Of New Haven Research Scholarship.

References

- [1] Blair, D.E., *Riemannian geometry of contact and symplectic manifolds*, Birkhauser, Boston, 2010.
- [2] Blair, D.E., Koufogiorgos, T. and Papantoniou, B. J., Contact metric manifolds satisfying a nullity condition, *Israel J. Math.* 91 (1995), 189-214.
- [3] Blair, D.E. and Sharma, R., Generalization of Myers' theorem on a contact manifold, *Illinois J. Math.* 34 (1990), 385-390.
- [4] Bochner, S., Curvature and Betti numbers, II, *Annals of Math.* 50 (1949), 77-93.
- [5] Bryant, R.L., Bochner-Kaehler metrics, *Jour. Amer. Math. Soc.* 14 (2001), 623-715.
- [6] Chen, B.Y., Some topological obstructions to Bochner-Kaehler metrics and their applications, *J. Diff. Geom.* 13 (1978), 547-558.
- [7] Ghosh, A. and Sharma, R., Almost Hermitian manifolds admitting holomorphically planar conformal vector fields, *J. Geom.* 84 (2005), 45-54.

- [8] Goldberg, S.I., *Curvature and Homology*, Academic Press, N.Y., 1962.
- [9] Koufogiorgos, T., On a class of contact Riemannian 3-manifolds, *Results in Math.* 27 (1995), 51-62.
- [10] Koufogiorgos, T., Contact strongly pseudo-convex integrable CR metrics as critical points, *J. Geom.* 59 (1997), 94-102.
- [11] Koufogiorgos, T., and Stamatiou, G., Strongly locally φ -symmetric contact metric manifolds, *Beitr. Algebra Geom.* 52 (2011), 221-236.
- [12] Obata, M., Certain conditions for a Riemannian manifold to be isometric with a sphere, *J. Math. Soc. Japan* 14 (1962), 333-340.
- [13] Okumura, M., Contact hypersurfaces in certain Kaehlerian manifolds, *Tohoku Math. J.* 18 (1966), 74-102.
- [14] Okumura, M., On infinitesimal conformal and projective transformations of normal contact spaces, *Tohoku Math. J.* 14 (1962), 398-412.
- [15] Sharma, R., Contact hypersurfaces of Kaehler manifolds, *J. Geom.* 78 (2003), 157-167.
- [16] Tanno, S., Variational problems on contact Riemannian manifolds, *Trans. Amer. Math. Soc.* 314(1989), 349-379.
- [17] Webster, S.M., On the pseudo-conformal geometry of a Kaehler manifold, *Math. Z.* 157 (1977), 265-270.
- [18] Yano, K., *Integral formulas in Riemannian geometry*, Marcel Dekker, New York, 1970.
- [19] Yano, K. and Kon, M., *Structures on Manifolds*, Series in Pure Mathematics 3, World Scientific Pub. Co., Singapore, 1984.