



University of
New Haven

University of New Haven

Digital Commons @ New Haven

Forensic Science Publications

Forensic Science

1-2020

Potential Applications of MicroRNA Profiling to Forensic Investigations

Claire L. Glynn

University of New Haven, cglynn@newhaven.edu

Follow this and additional works at: <https://digitalcommons.newhaven.edu/forensicscience-facpubs>



Part of the [Forensic Science and Technology Commons](#)

Publisher Citation

REVIEW: Potential applications of microRNA profiling to forensic investigations Claire L. Glynn RNA January 2020 26: 1-9; Published in Advance October 28, 2019, doi:10.1261/rna.072173.119

Comments

© 2020 Glynn; Published by Cold Spring Harbor Laboratory Press for the RNA Society

This article is distributed exclusively by the RNA Society for the first 12 months after the full-issue publication date (see <http://rnajournal.cshlp.org/site/misc/terms.xhtml>). After 12 months, it is available under a Creative Commons License (Attribution-NonCommercial 4.0 International), as described at <http://creativecommons.org/licenses/by-nc/4.0/>. This license permits non-commercial use, including reproduction, adaptation, and distribution of the article provided the original author and source are credited.

Article is online at <http://www.rnajournal.org/cgi/doi/10.1261/rna.072173.119>.

REVIEW

Potential applications of microRNA profiling to forensic investigations

CLAIRE L. GLYNN

Department of Forensic Science, Henry C. Lee College of Criminal Justice and Forensic Sciences, University of New Haven, West Haven, Connecticut 06516, USA

ABSTRACT

Within the forensic science community, there is a continued push to develop novel tools to aid in criminal investigations. microRNA (miRNA) analysis has been the focus of many researcher's attention in the biomedical field since its discovery in 1993; however, the forensic application of miRNA analysis has only been suggested within the last 10 years and has been gaining considerable traction recently. The primary focus of the forensic application of miRNA analysis has been on body fluid identification to provide confirmatory universal analysis of unknown biological stains obtained from crime scenes or evidence items. There are, however, other forensic applications of miRNA profiling that have shown potential, yet are largely understudied, and warrant further investigation such as organ tissue identification, donor age estimation, and more. This review paper aims to evaluate the current literature and future potential of miRNA analysis within the forensic science field.

Keywords: microRNA profiling; forensic science; body fluid identification; forensic investigations; organ tissue source identification

INTRODUCTION

microRNAs (miRNAs) are short, single-stranded, noncoding RNA molecules, typically 20–24 nt in length. The study of miRNAs began with the finding of their functional ability to regulate protein production in 1993 with the discovery of lin-4 by Lee et al. (1993). Seven years later, the second miRNA was discovered, let-7 (Pasquinelli et al. 2000), and miRNAs were subsequently established as a new class of riboregulators (Lagos-Quintana et al. 2001; Lau et al. 2001; Lee and Ambros 2001). Within the biomedical research community, miRNA analysis truly came to fruition when it was discovered that miRNAs play critical roles in a wide variety of biological and pathological processes, and that miRNAs were actually tissue-specific and could provide a signature of disease (Calin et al. 2002; Ambros 2004; Bartel 2004; Lu et al. 2005; Etheridge et al. 2011; Wegman and Krylov 2013). Since then there has been an explosion of interest in miRNAs and their diverse and wide-ranging capabilities within many disciplines of biological research. This research has mainly focused on clinical biological samples, such as various tissues (cancer vs. normal), cerebrospinal fluids, and cultured primary tissues. Comparatively, minimal research has focused on forensically relevant samples (e.g., body fluids, organ tissues, trace

deposits). The first suggestion of the forensic application of miRNAs was in 2009, when Hanson et al. (2009) suggested their potential for forensic body fluid identification (BFID). In the decade since, several researchers worldwide have investigated the potential for the forensic application of miRNAs. Despite the great promise miRNAs have shown in the last decade in terms of their forensic application, this field still remains very much in its infancy and has not yet been implemented in active forensic investigations or operational crime laboratory protocols. With the advancement of molecular methodologies and the sophistication of the techniques now used, miRNA profiling represents an ideal tool to supplement the forensic scientist's arsenal. Further studies and validation, however, are very much warranted, and required, in order to truly reveal the forensic potential of miRNA profiling, and certainly to withstand the scrutiny necessary for legal presentation in the courtroom.

BODY FLUID IDENTIFICATION

The identification of forensically relevant body fluids in a forensic investigation can provide important and probative

Corresponding author: cglynn@newhaven.edu

Article is online at <http://www.majournal.org/cgi/doi/10.1261/rna.072173.119>.

© 2020 Glynn This article is distributed exclusively by the RNA Society for the first 12 months after the full-issue publication date (see <http://majournal.cshlp.org/site/misc/terms.xhtml>). After 12 months, it is available under a Creative Commons License (Attribution-NonCommercial 4.0 International), as described at <http://creativecommons.org/licenses/by-nc/4.0/>.

evidence. Unambiguously confirming the body fluid source of a DNA profile is critical in forensic investigations as it clarifies the circumstances of an act, aids in determining the sequence of events, confirms or refutes victims/suspects/witness statements, and ultimately can provide linkages between victim/perpetrator/scene. Although there are both presumptive and confirmatory tests in routine practice for the commonly encountered forensically relevant body fluids (e.g., blood, semen, saliva, and urine), there are numerous limitations to these tests. Current BFID tests are based on chemical, enzymatic, or immunological reactions and typically have good sensitivity. However, their specificity is limited, with many tests presenting cross-reactivity with other body fluids, and they also produce a number of false positives (Virkler and Lednev 2009). Current tests, for the most part, are destructive to a sample, which is a major concern when dealing with already trace amounts of sample, and sufficient sample is required for subsequent DNA profiling to identify the donor of the body fluid. In addition, there are some body fluids, such as vaginal material and menstrual blood, which are greatly lacking in widely accepted protocols for their identification. Further to this, there is a great need for the development of a universal body fluid test, thereby eliminating the requirement to perform multiple individual body fluid tests, and having, instead, one test for them all.

Last, in an effort to streamline forensic biology work processes within operational crime laboratories, there is a need for the development of a method that also simultaneously contributes to the production of a DNA profile of the donor of the body fluid. Therefore, a method in which the sample preparation required for the body fluid test also provides DNA content for profiling would greatly improve the efficiency of the testing (see Fig. 1). In essence, the “ideal” forensic BFID test would be sensitive, specific, nondestructive, universal, and efficient.

Currently, there are significant efforts in the forensic science research field to develop novel methods for the identification of body fluids, to address the challenges currently faced. Previously, messenger RNA (mRNA) was thought to be a suitable candidate for the identification of body fluids. However, it was subsequently shown that mRNAs are susceptible to degradation and therefore are not always suitable for all forensically relevant samples—in particular, those samples that have been exposed to harsh environments (Hall and Ballantyne 2004; Setzer et al. 2008; Vennemann and Koppelkamm 2010; Haas et al. 2013; Hall et al. 2014). miRNAs, however, contain a wealth of information and are significantly more robust (Lee et al. 1993; Ambros 2001; Winter et al. 2009) than their mRNA counterparts, because of their smaller size. The tissue specificity of miRNAs discovered in the biomedical field




FIGURE 1. Workflow for universal microRNA body fluid identification with simultaneous DNA profiling.

sparked the idea that miRNAs could be utilized as markers for body fluids in forensic investigations (Hanson et al. 2009; Courts and Madea 2010), whereby specific body fluids would have specific miRNA signatures/expression profiles. Furthermore, forensic samples are often minute traces, and therefore any test that consumes the sample poses the risk of not having enough sample remaining after testing in order to obtain a DNA profile. The total nucleic acid content, however, can be extracted from a sample by means of a coextraction, producing two fractions: the DNA content and the total RNA component. This allows efficient processing of forensic samples, with BFID and DNA profiling occurring in parallel (Bauer and Patzelt 2003; Alvarez et al. 2004; Bowden et al. 2011; van der Meer et al. 2013; Li et al. 2014; Watanabe et al. 2014). Further to this, a recent study reports the ability to detect miRNAs in DNA extracts obtained using commonly used forensic DNA extraction methods (e.g., QIAamp DNA Investigator Kit [QIAGEN]) without any modifications to the protocols (Lewis et al. 2019). This highlights the ease of implementation of miRNA profiling into current crime laboratory workflows, whereby separate RNA extraction or coextraction methods are not strictly necessary. Moreover, potential exists for miRNA profiling to be performed in cold case investigations, in which only DNA extracts remain from the original evidence item submitted.

Previous published studies reported promising results identifying differentially expressed miRNAs with great potential as novel biomarkers for forensic BFID (Hanson et al. 2009; Zubakov et al. 2010; Courts and Madea 2011; Wang et al. 2013; Sauer et al. 2016; Seashols-Williams et al. 2016;

Sirker et al. 2017; Mayes et al. 2018; O'Leary and Glynn 2018; Tian et al. 2018). In the small number of studies published in this area, however, there is little agreement between them, with only some miRNAs suggested as potential biomarkers for specific fluids overlapping between studies (see Table 1).

The first to explore the forensic application of miRNAs was Erin K. Hanson in 2009 (Hanson et al. 2009). At first, researchers aimed to discover miRNAs that are specific to one particular body fluid; however, research studies have been unsuccessful in this search thus far (Hanson et al. 2009; Zubakov et al. 2010; Courts and Madea 2011; Wang et al. 2013; Sauer et al. 2016; Seashols-Williams et al. 2016; Sirker et al. 2017; Mayes et al. 2018; Tian et al. 2018). However, Hanson et al. (2009) revealed nine miRNAs that were sufficiently differentially expressed to such a degree as to permit the identification of the particular body fluid. In addition, Hanson et al. (2009) revealed that as little as 50 pg of total RNA was needed, highlighting its utility with minute forensic samples. Zubakov et al. (2010) screened a large set of 718 human miRNA markers in forensically relevant body fluids, using the most comprehensive microarray platform available at that time. The results identified a number of candidate miRNAs for each body fluid. However, upon further validation using reverse transcription quantitative polymerase chain reaction (RT-qPCR), only two miRNAs for blood and two for semen were suggested as suitable for future forensic applications. Furthermore, this study also revealed the time-wise stability of miRNAs over 1 year in ambient laboratory conditions and also the sensitivity of this technique, as reliable marker

TABLE 1. miRNAs identified in the literature as showing potential as a biomarker for individual body fluids

Body fluid	Mayes et al. (2018)	Sirker et al. (2017)	Seashols-Williams et al. (2016)	Sauer et al. (2016)	Wang et al. (2013)	Courts and Madea (2011)	Zubakov et al. (2010)	Hanson et al. (2009)
Venous blood	miR-142-3p, miR-451	miR-16, miR-451	miR-200b	miR-144-3p, miR-144-5p, miR-451a	miR-16, miR-486	miR-126, miR-150, miR-451	miR-20a, miR-106a, miR-185, miR-144	miR-16, miR-451
Menstrual blood	miR-141-3p, miR-412-3p	miR-185, miR-412-3p	miR-200b	miR-144-3p, miR-144-5p, miR-451a	miR-214	–	miR-144, miR-185	miR-412, miR-451
Semen	miR-10b, miR-891a	miR-10b, miR-943	miR-891a	miR-10a, miR-10b, miR-135a	miR-888, miR-891a	miR-200, miR-203, miR-205	miR-10a, miR-135a, miR-507, miR-943, miR-891a	miR-10b, miR-135b
Saliva	miR-205	miR-124, miR-203, miR-205	miR-26b	–	miR-138	–	miR-208b, miR-518c, miR-583	miR-205, miR-658
Vaginal material	–	miR-1280, miR-4286	–	–	–	–	miR-617, miR-891a	miR-124a, miR-891a

(–) Not studied/none found.

detection was obtained with just picograms of starting material.

Wang et al. (2013) published a table illustrating some of the previous miRNA findings for particular body fluids in their research and other groups. Wang et al. performed an array that examined 754 human-specific miRNAs. Currently, the total count of discovered miRNAs that identify with *Homo sapiens* is ~2700 according to the miRBase registry (release 22, accessed March 2019). Another database/registry, called MirGeneDB, compiled using primary high-throughput sequencing data, currently lists only 556 authentic human miRNA genes (Fromm et al. 2015). It could be suggested that perhaps a truly body fluid-specific miRNA has simply not yet been realized; however, by comparing both registries and considering only the authentic miRNAs, it appears that only a small number of new human miRNAs have been identified in the last decade. Therefore, it is unlikely that a new truly body fluid-specific miRNA that would have sufficient expression to enable quantification will be realized.

Although the discovery of truly specific miRNAs to individual body fluids would be of great benefit, it is reasonable to assume that the differential expression of multiple miRNA targets would provide BFID with sufficient accuracy. Sauer et al. (2016) published a comprehensive study using a thoroughly validated, state-of-the-art qPCR procedure and reported four miRNAs that were shown to be useful for differentiation. This study expanded further by using a decision algorithm to detect each of the five body fluids by using as few markers as possible to simplify the analysis procedure. The results, however, were not as discriminatory as expected, highlighting the fact that a panel of miRNAs for each body fluid is truly needed. RT-qPCR is currently the method of choice for differential expression analysis of individual miRNA targets. However, to ensure reliable detection and expression analysis of miRNAs for forensic BFID, it is essential that proper normalization strategies using endogenous controls are used. Sirker et al. (2017) identified four miRNAs as the most stably expressed across the set of samples and that, therefore, could act as potential endogenous controls. Further to this, five miRNAs were reported to be successful in the differentiation of six different cell types/body fluids, using the previously identified endogenous controls to normalize the qPCR data. However, this study was not able to differentiate between venous and menstrual blood, which would be preferable for forensic applications. This study further highlighted the lack of exclusive marker specificity for one body fluid and a simple interpretation method. Both Sauer et al. (2016) and Sirker et al. (2017) stressed the importance, and benefits of, interlaboratory trials using a set of ubiquitous markers to better assess the experimental approach and ultimately its suitability for forensic casework implementation.

Seashols-Williams et al. (2016) published the first report of forensically relevant body fluids subjected to next-generation sequencing (NGS) using the Illumina Hi-Seq platform. Also, Wang et al. (2016) reported small RNA sequencing of blood and saliva using the Ion Personal Genome Machine System (Wang et al. 2016). Seashols-Williams et al. (2016) provided a very comprehensive and well-designed study that examined every step in the workflow. The authors identified a suitable RNA extraction procedure for forensically relevant body fluids and suitable endogenous controls for normalization in RT-qPCR, and they produced a large data set, with the consequent development of candidate miRNAs for further research. More recently, Dørum et al. (2019) applied whole miRNome massively parallel sequencing to six forensically relevant body fluids. Their findings were in agreement with those of Seashols-Williams et al., in particular with regard to the reference miRNA markers.

Once panels of miRNAs markers are identified and validated for BFID, it will be crucial to apply appropriate statistical methods to the data analysis. Validated multivariate statistical methods are certainly required for the analysis of differentially expressed miRNAs using RT-qPCR. The $\Delta\text{CT}/\Delta\Delta\text{CT}$ methods are typically used in relative quantification (RQ) studies. Many of these studies report using one-way ANOVA and independent two-sample *t*-tests to determine association and comparisons between individual body fluids and to assess significance (Courts and Madea 2011; Hanson and Ballantyne 2013; Li et al. 2017). Dørum et al. (2019) reported using partial least squares (PLS) and linear discriminant analysis (LDA) to predict body fluids and also identified a minimum number of miRNA markers that will still provide good prediction accuracy. Other statistical approaches have been applied, such as that of Hanson et al. (2014), who developed a quantitative statistical model using logistic regression to predict menstrual blood and reported a high, and measurable, degree of accuracy.

Although they are few, the previous research studies have highlighted the potential of miRNA markers for the identification of forensically relevant body fluids. However, further research is required to identify panels of unambiguous markers for each of the forensically relevant body fluids. Although the research studies discussed here contribute greatly to the growing body of knowledge on this topic, there is yet more research to be performed, however, and it must be assessed if results previously obtained are reproducible using different platforms and approaches. Further to this, some forensically relevant body fluids have been less extensively explored (e.g., vaginal fluid and menstrual blood). With further research, the potential of miRNAs for forensic BFID continues to grow. Should this novel method one day be implemented into routine casework, it will address current challenges encountered in body fluid testing by creating a sensitive,

specific, universal test for BFID, which will also allow for the coextraction of both the DNA and RNA content in one process, thereby streamlining workflow processes and ultimately forensic practice.

ORGAN TISSUE IDENTIFICATION

Similar to BFID, the identification of certain internal organ tissues can yield valuable probative evidence in the investigation of a violent crime. During a violent act, internal organ tissue may be transferred from the victim to the perpetrator, to the weapon or innocuous item present, or to the location/scene at which the crime is being committed. Internal organ tissues may adhere to a bullet passing through a body or to a knife that has penetrated the skin. In certain circumstances, identifying the organ tissue source of a DNA profile obtained from an evidence item, such as a knife blade or a bullet, confirms the role that evidence item played in the crime being investigated. Furthermore, the ability to identify organ tissues in trace deposits of biological material located at scenes (e.g., in the plumbing or sewers of a building) can provide investigative leads to the possibility that the human remains have been removed to another location or the dismemberment or disposition of human remains at that location. Currently, there are no standard molecular operating procedures for organ tissue identification used in routine casework at operational crime laboratories. Investigators are limited to simply identifying the DNA profile of the donor of the specimen on the evidence item and can only suggest possible organ tissue sources of the DNA. If organ tissue identification was deemed crucial to an investigation, the sample would likely be sent out for cellular/histological analysis to either the Office of the Chief Medical Examiner (OCME) (only in the United States), State Pathologist, or to an external biological analysis company, where immunohistochemical analyses would likely be performed. Other traditional methods, using immunological and enzymatic processes, have previously been suggested as viable tools for this purpose, although none ever gained traction within the forensic science industry (Kimura et al. 1995; Takahama 1996; Seo et al. 1997; Takata et al. 2004). The most confounding limitation of all of the traditional methods is that they require a relatively large sample size that is in near-pristine condition. In forensic investigations, specimens are often minute in size and heavily degraded; therefore, these traditional methods are not ideally suited for forensic use.

In recent years, more molecular-based methods, such as mRNA and DNA methylation analysis, for organ tissue identification have been suggested as viable tools, with some researchers producing promising results (Hanson and Ballantyne 2017; van den Berge and Sijen 2017; Samsuwan et al. 2018). As miRNAs have shown great

promise for BFID, it is logical to infer that same potential to their use for organ tissue identification, in particular because of the miRNA's robustness and ability to withstand degradation. Sauer et al. (2017) published the first report of the use of miRNA markers for the identification of brain, kidney, liver, lung, skin, heart muscle, and skeletal muscle. The researchers provided a comprehensive investigation by performing RT-qPCR of 15 preselected miRNAs on not just pristine samples, but also mixed, aged, and degraded samples. Further to this, the researchers generated samples from mock stabbings and shootings to ensure the robustness of this method to realistic forensic specimens. Although the study identified individual miRNAs that were differentially expressed and therefore showed potential as novel biomarkers for individual organ tissues, it was noted that multiple miRNAs or panels of miRNAs are advisable for the inference of certain tissues. The authors also note that further research should comprise larger sample sets and an expansion of the organs investigated, even suggesting organ subsections. However, this study provides a strong foundation from which further research into this novel application of miRNAs can now be explored.

OTHER POTENTIAL FUTURE APPLICATIONS

In the biomedical field, human aging is an area of great interest, in particular with regard to identifying diagnostic predictors of age-associated diseases, such as cancer. There is potential to translate the findings of biomedical research studies to forensic applications for estimating the age of a donor of a particular biological sample using miRNA analysis. Noren Hooten et al. (2010) profiled 800 miRNAs in both young (~30 yr) and old (~64 yr) cohorts, and reported that the majority of miRNAs decreased in abundance with age. This suggests that changes in miRNA expression could have potential as an indicator of the age of the donor. Similarly, determining the age of the actual stain—or as it is more commonly known, the time since deposition (TSD)—is often questioned in forensic investigations, as it could relate to the stain's relevance to the crime and also to determining the sequence of events. mRNAs were previously investigated as a potential tool for this purpose (Anderson et al. 2005); however, the wide range and instability of mRNA suggests this may not be an effective tool. One study reported three miRNAs (Mohammed et al. 2018) to steadily and significantly increase over time, whereas another (Alshehhi and Hadrill 2019) investigated the use of both mRNA and miRNA markers and reported that each marker exhibited a unique degradation profile over a 1-yr time period. Although these results are promising, further research is certainly required to assess the impact of environmental insults. Last, postmortem interval (PMI) estimation has recently been suggested as a novel application of miRNA

analysis. PMI is defined as the period of time elapsed from the time of death. Currently, a variety of methods are used to estimate PMI (e.g., physical, physiochemical, biochemical, microbiological, entomological, and botanical processes) (Henssge and Madea 2007); however, none are absolute. The degradation of nucleic acids (both DNA and RNA) have been well-studied, and it recently has been suggested that nucleic acids could be ideal biomarkers of PMI (Di Nunno et al. 1998; Bauer et al. 2003; Liu et al. 2007; Sampaio-Silva et al. 2013; Scrivano et al. 2019). Both miRNAs and circulating RNAs (circRNAs) have shown satisfactory stability across a range of tissue types in the post-mortem interval, highlighting that miRNAs are less susceptible to their environmental conditions and therefore suitable as stable reference genes (Tu et al. 2019). Although the results of this research are promising, there are many variables to be considered in estimating PMI, which are much more complex in real-life scenarios with humans than in laboratory conditions using animal models.

POTENTIAL LIMITATIONS

Should miRNA profiling be implemented into routine practice within forensic investigations, it is crucial that it is robust enough to handle the challenges that forensic samples typically bring (e.g., nonpristine environmentally damaged samples). Proteins, as well as nucleic acids, have all been found to degrade, lose conformation, and, subsequently, lose function over time. These processes are shown to be accelerated when exposed to various environmental insults (Pfeifer et al. 2005; Alaeddini et al. 2010). Some of these are, but not necessarily limited to, prolonged ultraviolet light exposure (Hall and Ballantyne 2004; Hall et al. 2014), excessive heat exposure (Lindahl and Nyberg 1974), water/moisture (Marrone and Ballantyne 2009, 2010), and various forms of chemical/enzymatic damage (Gates 2009). Various studies over the last decade have elucidated the nature of some of the damage mechanisms associated with biological evidence (Hall and Ballantyne 2004; Gates 2009; Marrone and Ballantyne 2009, 2010; Hall et al. 2014). DNA damage with regard to forensic specimens has been widely studied; various methods exist that attempt to repair various forms of DNA damage (Evans and Nichols 2008; Diegoli et al. 2012; Ambers et al. 2014; Robertson et al. 2014; Wallace 2014). Little is known, however, with regard to miRNA damage—in particular, with forensic specimens. There has been no published research to date that investigates the impact of damage on the RNA content of forensically relevant body fluids—more specifically the miRNA content—nor has there been any published research that explores the ability to “repair” the damaged RNA content of forensically relevant body fluids.

Uniformity and acceptance of methods across research studies has certainly presented limitations to date. The lack of agreement in the studies discussed previously (Table 1) could certainly be as a result of the variety of methods used across them. It is imperative that uniform and agreed-on methods are utilized on a global level, to ensure data obtained from different research groups or crime laboratories can be reliably compared. This includes agreement on the starting sample volume and concentration, the method of extraction and subsequent quantification, followed by the method of amplification and target analysis, and concluding with statistical/interpretation analysis to be performed. As miRNA research has progressed in the biomedical field, the RT-qPCR technique has become the gold standard for the analysis of individual miRNA targets, because of its ease of use and the robustness of the technique (Ach et al. 2008; Zubakov et al. 2010; Omelia et al. 2013; Sauer et al. 2016). NGS, however, has recently gained widespread attention for miRNA analysis. NGS for small RNAs allows the sequencing of small RNA species with unprecedented sensitivity and dynamic range. It is possible to discover novel miRNAs and also examine differential expression of all small RNAs in a sample. It has been suggested that NGS platforms will be standard instrumentation in crime laboratories in the future, as the advantages are many, in addition to their modes of applications within the forensic science discipline. As a result, all available NGS platforms must be investigated and compared to ensure the most appropriate platform is implemented. As stated previously, the ultimate goal is to identify panels of miRNAs for each individual body fluid to be used for identification; therefore, RT-qPCR may not be the ideal approach as multiple multiplexes and consumption of the sample would be required. NGS, although costly, would provide a more comprehensive, yet streamlined approach, because of its multiplexing nature and the ease of interrogating the whole or a large subset of the miRNAome.

Last, it is the author’s opinion that there are three further reasons that contribute to keeping the forensic application of miRNAs from progressing to the level it deserves. First, there is a limited amount of funding provided for forensic science research applications, with novel research associated with forensic DNA profiling for person identification purposes often taking precedence over RNA profiling applications. Second, the required expertise in the area of miRNA analysis is predominantly focused in the biomedical/clinical field. Until sufficient funding is provided to the RNA-based forensic science discipline, however, it remains difficult to attract talented researchers to contribute to fields of study such as this. Last, the lack of commercially available kits and reagents designed specifically for forensic use is a further impediment to the implementation of miRNA and other RNA technologies in forensic casework applications.

CONCLUSION

Although microRNA expression analysis is being widely explored the world over, this exploration has focused predominantly on its application to the biomedical/cancer research field. To truly reveal the full potential of miRNAs within the forensic field, it is crucial to unravel the molecular complexities of the samples at hand. The implementation of miRNA analysis into the forensic workflow as described above would serve to address several challenges currently faced by forensic scientists. By being able to establish robust performance under adverse conditions it allows for the reliable access to information that was not previously possible. It is time to move the state of the art forward, by addressing current challenges with new tools, thus allowing forensic scientists to harvest more layers of information from biological evidence. As mentioned previously, the forensic application of microRNA analysis is only just being realized. As a result, published research in this area is limited, yet growing. With the diverse array of forensic applications for miRNA analysis it is possible to pioneer novel approaches to physical evidence. In forensic science, methods that once seemed state-of-the-art, now are becoming antiquated. It is crucial that researchers work toward developing techniques for use in forensic investigations that produce results in record time, providing higher powers of discrimination, and that are sufficiently robust to gain acceptance by both the scientific community and, of course, the courtroom.

REFERENCES

- Ach RA, Wang H, Curry B. 2008. Measuring microRNAs: comparisons of microarray and quantitative PCR measurements, and of different total RNA prep methods. *BMC Biotechnol* **8**: 69. doi:10.1186/1472-6750-8-69
- Alaeddini R, Walsh SJ, Abbas A. 2010. Forensic implications of genetic analyses from degraded DN—a review. *Forensic Sci Int Genet* **4**: 148–157. doi:10.1016/j.fsigen.2009.09.007
- Alshehhi S, Haddrill PR. 2019. Estimating time since deposition using quantification of RNA degradation in body fluid—specific markers. *Forensic Sci Int* **298**: 58–63. doi:10.1016/j.forsciint.2019.02.046
- Alvarez M, Juusola J, Ballantyne J. 2004. An mRNA and DNA co-isolation method for forensic casework samples. *Anal Biochem* **335**: 289–298. doi:10.1016/j.ab.2004.09.002
- Ambers A, Turnbough M, Benjamin R, King J, Budowle B. 2014. Assessment of the role of DNA repair in damaged forensic samples. *Int J Legal Med* **128**: 913–921. doi:10.1007/s00414-014-1003-3
- Ambros V. 2001. microRNAs: tiny regulators with great potential. *Cell* **107**: 823–826. doi:10.1016/S0092-8674(01)00616-X
- Ambros V. 2004. The functions of animal microRNAs. *Nature* **431**: 350–355. doi:10.1038/nature02871
- Anderson S, Howard B, Hobbs GR, Bishop CP. 2005. A method for determining the age of a bloodstain. *Forensic Sci Int* **148**: 37–45. doi:10.1016/j.forsciint.2004.04.071
- Bartel DP. 2004. MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell* **116**: 281–297. doi:10.1016/S0092-8674(04)00045-5
- Bauer M, Patzelt D. 2003. A method for simultaneous RNA and DNA isolation from dried blood and semen stains. *Forensic Sci Int* **136**: 76–78. doi:10.1016/S0379-0738(03)00219-6
- Bauer M, Gramlich I, Polzin S, Patzelt D. 2003. Quantification of mRNA degradation as possible indicator of postmortem interval—a pilot study. *Leg Med (Tokyo)* **5**: 220–227. doi:10.1016/j.legalmed.2003.08.001
- Bowden A, Fleming R, Harbison S. 2011. A method for DNA and RNA co-extraction for use on forensic samples using the Promega DNA IQ system. *Forensic Sci Int Genet* **5**: 64–68. doi:10.1016/j.fsigen.2009.11.007
- Calin GA, Dumitru CD, Shimizu M, Bichi R, Zupo S, Noch E, Aldler H, Rattan S, Keating M, Rai K, et al. 2002. Frequent deletions and down-regulation of micro-RNA genes *miR15* and *miR16* at 13q14 in chronic lymphocytic leukemia. *Proc Natl Acad Sci* **99**: 15524–15529. doi:10.1073/pnas.242606799
- Courts C, Madea B. 2010. Micro-RNA—a potential for forensic science? *Forensic Sci Int* **203**: 106–111. doi:10.1016/j.forsciint.2010.07.002
- Courts C, Madea B. 2011. Specific micro-RNA signatures for the detection of saliva and blood in forensic body-fluid identification. *J Forensic Sci* **56**: 1464–1470. doi:10.1111/j.1556-4029.2011.01894.x
- Diegoli TM, Farr M, Cromartie C, Coble MD, Bille TW. 2012. An optimized protocol for forensic application of the PreCR Repair Mix to multiplex STR amplification of UV-damaged DNA. *Forensic Sci Int Genet* **6**: 498–503. doi:10.1016/j.fsigen.2011.09.003
- Di Nunno NR, Costantinides F, Bernasconi P, Bottin C, Melato M. 1998. Is flow cytometric evaluation of DNA degradation a reliable method to investigate the early postmortem period? *Am J Forensic Med Pathol* **19**: 50–53. doi:10.1097/00000433-199803000-00008
- Dørum G, Ingold S, Hanson E, Ballantyne J, Russo G, Aluri S, Snipen L, Haas C. 2019. Predicting the origin of stains from whole miRNome massively parallel sequencing data. *Forensic Sci Int Genet* **40**: 131–139. doi:10.1016/j.fsigen.2019.02.015
- Etheridge A, Lee I, Hood L, Galas D, Wang K. 2011. Extracellular microRNA: a new source of biomarkers. *Mutat Res* **717**: 85–90. doi:10.1016/j.mrfmmm.2011.03.004
- Evans TC, Nichols NM. 2008. DNA repair enzymes. *Curr Protoc Mol Biol* **84**: 3.9.1–3.9.12. doi:10.1002/0471142727.mb0309s84
- Fromm B, Billipp T, Peck LE, Johansen M, Tarver JE, King BL, Newcomb JM, Sempere LF, Flatmark K, Hovig E, et al. 2015. A uniform system for the annotation of vertebrate microRNA genes and the evolution of the human microRNAome. *Annu Rev Genet* **49**: 213–242. doi:10.1146/annurev-genet-120213-092023
- Gates KS. 2009. An overview of chemical processes that damage cellular DNA: spontaneous hydrolysis, alkylation, and reactions with radicals. *Chem Res Toxicol* **22**: 1747–1760. doi:10.1021/tx900242k
- Haas C, Hanson E, Ballantyne J. 2013. mRNA and microRNA for body fluid identification. In *Encyclopedia of forensic sciences*, pp. 402–408.
- Hall A, Ballantyne J. 2004. Characterization of UVC-induced DNA damage in bloodstains: forensic implications. *Anal Bioanal Chem* **380**: 72–83. doi:10.1007/s00216-004-2681-3
- Hall A, Sims LM, Ballantyne J. 2014. Assessment of DNA damage induced by terrestrial UV irradiation of dried bloodstains: forensic implications. *Forensic Sci Int Genet* **8**: 24–32. doi:10.1016/j.fsigen.2013.06.010
- Hanson EK, Ballantyne J. 2013. Circulating microRNA for the identification of forensically relevant body fluids. *Methods Mol Biol* **1024**: 221–234. doi:10.1007/978-1-62703-453-1_18
- Hanson E, Ballantyne J. 2017. Human organ tissue identification by targeted RNA deep sequencing to aid the investigation

- of traumatic injury. *Genes (Basel)* **8**: 319. doi:10.3390/genes8110319
- Hanson EK, Lubenow H, Ballantyne J. 2009. Identification of forensically relevant body fluids using a panel of differentially expressed microRNAs. *Anal Biochem* **387**: 303–314. doi:10.1016/j.ab.2009.01.037
- Hanson EK, Mirza M, Rehak K, Ballantyne J. 2014. The identification of menstrual blood in forensic samples by logistic regression modeling of miRNA expression. *Electrophoresis* **35**: 3087–3095. doi:10.1002/elps.201400171
- Hensge C, Madea B. 2007. Estimation of the time since death. *Forensic Sci Int* **165**: 182–184. doi:10.1016/j.forsciint.2006.05.017
- Kimura A, Ikeda H, Yasuda S, Yamaguchi K, Tsuji T. 1995. Brain tissue identification based on myosin heavy chain isoforms. *Int J Legal Med* **107**: 193–196. doi:10.1007/BF01428404
- Lagos-Quintana M, Rauhut R, Lendeckel W, Tuschl T. 2001. Identification of novel genes coding for small expressed RNAs. *Science* **294**: 853–858. doi:10.1126/science.1064921
- Lau NC, Lim LP, Weinstein EG, Bartel DP. 2001. An abundant class of tiny RNAs with probable regulatory roles in *Caenorhabditis elegans*. *Science* **294**: 858–862. doi:10.1126/science.1065062
- Lee RC, Ambros V. 2001. An extensive class of small RNAs in *Caenorhabditis elegans*. *Science* **294**: 862–864. doi:10.1126/science.1065329
- Lee RC, Feinbaum RL, Ambros V. 1993. The *C. elegans* heterochronic gene *lin-4* encodes small RNAs with antisense complementarity to *lin-14*. *Cell* **75**: 843–854. doi:10.1016/0092-8674(93)90529-Y
- Lewis CA, Layne TR, Seashols-Williams SJ. 2019. Detection of microRNAs in DNA extractions for forensic biological source identification. *J Forensic Sci* **64**: 1823–1830. doi:10.1111/1556-4029.14070
- Li Y, Zhang J, Wei W, Wang Z, Prinz M, Hou Y. 2014. A strategy for co-analysis of microRNAs and DNA. *Forensic Sci Int Genet* **12**: 24–29. doi:10.1016/j.fsigen.2014.04.011
- Li Z, Bai P, Peng D, Wang H, Guo Y, Jiang Y, He W, Tian H, Yang Y, Huang Y, et al. 2017. Screening and confirmation of microRNA markers for distinguishing between menstrual and peripheral blood. *Forensic Sci Int Genet* **30**: 24–33. doi:10.1016/j.fsigen.2017.05.012
- Lindahl T, Nyberg B. 1974. Heat-induced deamination of cytosine residues in deoxyribonucleic acid. *Biochemistry* **13**: 3405–3410. doi:10.1021/bi00713a035
- Liu L, Shu X, Ren L, Zhou H, Li Y, Liu W, Zhu C, Liu L. 2007. Determination of the early time of death by computerized image analysis of DNA degradation: which is the best quantitative indicator of DNA degradation? *J Huazhong Univ Sci Technolog Med Sci* **27**: 362–366. doi:10.1007/s11596-007-0404-7
- Lu J, Getz G, Miska EA, Alvarez-Saavedra E, Lamb J, Peck D, Sweet-Cordero A, Ebert BL, Mak RH, Ferrando AA, et al. 2005. MicroRNA expression profiles classify human cancers. *Nature* **435**: 834–838. doi:10.1038/nature03702
- Marrone A, Ballantyne J. 2009. Changes in dry state hemoglobin over time do not increase the potential for oxidative DNA damage in dried blood. *PLoS One* **4**: e5110. doi:10.1371/journal.pone.0005110
- Marrone A, Ballantyne J. 2010. Hydrolysis of DNA and its molecular components in the dry state. *Forensic Sci Int Genet* **4**: 168–177. doi:10.1016/j.fsigen.2009.08.007
- Mayes C, Seashols-Williams S, Hughes-Stamm S. 2018. A capillary electrophoresis method for identifying forensically relevant body fluids using miRNAs. *Leg Med (Tokyo)* **30**: 1–4. doi:10.1016/j.legalmed.2017.10.013
- Mohammed AT, Khalil SR, Ali HA, Awad A. 2018. Validation of mRNA and microRNA profiling as tools in qPCR for estimation of the age of bloodstains. *Life Sci J* **15**: 1–7.
- Noren Hooten N, Abdelmohsen K, Gorospe M, Ejiogu N, Zonderman AB, Evans MK. 2010. microRNA expression patterns reveal differential expression of target genes with age. *PLoS One* **5**: e10724. doi:10.1371/journal.pone.0010724
- O’Leary KR, Glynn CL. 2018. Investigating the isolation and amplification of microRNAs for forensic body fluid identification. *MicroRNA* **7**: 187–194. doi:10.2174/2211536607666180430153821
- Omelia EJ, Uchimoto ML, Williams G. 2013. Quantitative PCR analysis of blood- and saliva-specific microRNA markers following solid-phase DNA extraction. *Anal Biochem* **435**: 120–122. doi:10.1016/j.ab.2012.12.024
- Pasquinelli AE, Reinhart BJ, Slack F, Martindale MQ, Kuroda MI, Maller B, Hayward DC, Ball EE, Degnan B, Muller P, et al. 2000. Conservation of the sequence and temporal expression of let-7 heterochronic regulatory RNA. *Nature* **408**: 86–89. doi:10.1038/35040556
- Pfeifer GP, You YH, Besaratinia A. 2005. Mutations induced by ultraviolet light. *Mutat Res* **571**: 19–31. doi:10.1016/j.mrfmmm.2004.06.057
- Robertson JM, Dineen SM, Scott KA, Lucyshyn J, Saeed M, Murphy DL, Schweighardt AJ, Meiklejohn KA. 2014. Assessing PreCR repair enzymes for restoration of STR profiles from artificially degraded DNA for human identification. *Forensic Sci Int Genet* **12**: 168–180. doi:10.1016/j.fsigen.2014.05.011
- Sampaio-Silva F, Magalhães T, Carvalho F, Dinis-Oliveira RJ, Silvestre R. 2013. Profiling of RNA degradation for estimation of post mortem [sic] interval. *PLoS One* **8**: e56507. doi:10.1371/journal.pone.0056507
- Samsuwan J, Muangsub T, Yanatatsaneejit P, Mutirangura A, Kitkumthorn N. 2018. Combined bisulfite restriction analysis for brain tissue identification. *Forensic Sci Int* **286**: 42–45. doi:10.1016/j.forsciint.2018.02.032
- Sauer E, Reinke AK, Courts C. 2016. Differentiation of five body fluids from forensic samples by expression analysis of four microRNAs using quantitative PCR. *Forensic Sci Int Genet* **22**: 89–99. doi:10.1016/j.fsigen.2016.01.018
- Sauer E, Extra A, Cachée P, Courts C. 2017. Identification of organ tissue types and skin from forensic samples by microRNA expression analysis. *Forensic Sci Int Genet* **28**: 99–110. doi:10.1016/j.fsigen.2017.02.002
- Scrivano S, Sanavio M, Tozzo P, Caenazzo L. 2019. Analysis of RNA in the estimation of post-mortem interval: a review of current evidence. *Int J Legal Med* **133**: 1629–1640. doi:10.1007/s00414-019-02125-x
- Seashols-Williams S, Lewis C, Calloway C, Peace N, Harrison A, Hayes-Nash C, Fleming S, Wu Q, Zehner ZE. 2016. High-throughput miRNA sequencing and identification of biomarkers for forensically relevant biological fluids. *Electrophoresis* **37**: 2780–2788. doi:10.1002/elps.201600258
- Seo Y, Kakizaki E, Takahama K. 1997. A sandwich enzyme immunoassay for brain S-100 protein and its forensic application. *Forensic Sci Int* **87**: 145–154. doi:10.1016/S0379-0738(97)00049-2
- Setzer M, Juusola J, Ballantyne J. 2008. Recovery and stability of RNA in vaginal swabs and blood, semen, and saliva stains. *J Forensic Sci* **53**: 296–305. doi:10.1111/j.1556-4029.2007.00652.x
- Sirker M, Fimmers R, Schneider PM, Gomes I. 2017. Evaluating the forensic application of 19 target microRNAs as biomarkers in body fluid and tissue identification. *Forensic Sci Int Genet* **27**: 41–49. doi:10.1016/j.fsigen.2016.11.012
- Takahama K. 1996. Forensic application of organ-specific antigens. *Forensic Sci Int* **80**: 63–69. doi:10.1016/0379-0738(96)01928-7
- Takata T, Miyaiishi S, Kitao T, Ishizu H. 2004. Identification of human brain from a tissue fragment by detection of neurofilament proteins. *Forensic Sci Int* **144**: 1–6. doi:10.1016/j.forsciint.2004.01.020

- Tian H, Lv M, Li Z, Peng D, Tan Y, Wang H, Li Q, Li F, Liang W. 2018. Semen-specific miRNAs: suitable for the distinction of infertile semen in the body fluid identification? *Forensic Sci Int Genet* **33**: 161–167. doi:10.1016/j.fsigen.2017.12.010
- Tu C, Du T, Ye X, Shao C, Xie J, Shen Y. 2019. Using miRNAs and circRNAs to estimate PMI in advanced stage. *Leg Med (Tokyo)* **38**: 51–57. doi:10.1016/j.legalmed.2019.04.002
- van den Berge M, Sijen T. 2017. Extended specificity studies of mRNA assays used to infer human organ tissues and body fluids. *Electrophoresis* **38**: 3155–3160. doi:10.1002/elps.201700241
- van der Meer D, Uchimoto ML, Williams G. 2013. Simultaneous analysis of micro-RNA and DNA for determining the body fluid origin of DNA profiles. *J Forensic Sci* **58**: 967–971. doi:10.1111/1556-4029.12160
- Vennemann M, Koppelkamm A. 2010. mRNA profiling in forensic genetics I: possibilities and limitations. *Forensic Sci Int* **203**: 71–75. doi:10.1016/j.forsciint.2010.07.006
- Virkler K, Lednev IK. 2009. Analysis of body fluids for forensic purposes: from laboratory testing to non-destructive rapid confirmatory identification at a crime scene. *Forensic Sci Int* **188**: 1–17. doi:10.1016/j.forsciint.2009.02.013
- Wallace SS. 2014. Base excision repair: a critical player in many games. *DNA Repair (Amst)* **19**: 14–26. doi:10.1016/j.dnarep.2014.03.030
- Wang Z, Zhang J, Luo H, Ye Y, Yan J, Hou Y. 2013. Screening and confirmation of microRNA markers for forensic body fluid identification. *Forensic Sci Int Genet* **7**: 116–123. doi:10.1016/j.fsigen.2012.07.006
- Wang Z, Zhou D, Cao Y, Hu Z, Zhang S, Bian Y, Hou Y, Li C. 2016. Characterization of microRNA expression profiles in blood and saliva using the Ion Personal Genome Machine[®] System (Ion PGM[™] System). *Forensic Sci Int Genet* **20**: 140–146. doi:10.1016/j.fsigen.2015.10.008
- Watanabe K, Iwashima Y, Akutsu T, Sekiguchi K, Sakurada K. 2014. Evaluation of a co-extraction method for real-time PCR-based body fluid identification and DNA typing. *Leg Med (Tokyo)* **16**: 56–59. doi:10.1016/j.legalmed.2013.11.002
- Wegman DW, Krylov SN. 2013. Direct miRNA-hybridization assays and their potential in diagnostics. *TrAC Trends Anal Chem* **44**: 121–130. doi:10.1016/j.trac.2012.10.014
- Winter J, Jung S, Keller S, Gregory RI, Diederichs S. 2009. Many roads to maturity: microRNA biogenesis pathways and their regulation. *Nat Cell Biol* **11**: 228–234. doi:10.1038/ncb0309-228
- Zubakov D, Boersma AW, Choi Y, van Kuijk PF, Wiemer EA, Kayser M. 2010. MicroRNA markers for forensic body fluid identification obtained from microarray screening and quantitative RT-PCR confirmation. *Int J Legal Med* **124**: 217–226. doi:10.1007/s00414-009-0402-3



RNA

A PUBLICATION OF THE RNA SOCIETY

Potential applications of microRNA profiling to forensic investigations

Claire L. Glynn

RNA 2020 26: 1-9 originally published online October 28, 2019
Access the most recent version at doi:[10.1261/rna.072173.119](https://doi.org/10.1261/rna.072173.119)

References

This article cites 74 articles, 4 of which can be accessed free at:
<http://rnajournal.cshlp.org/content/26/1/1.full.html#ref-list-1>

Creative Commons License

This article is distributed exclusively by the RNA Society for the first 12 months after the full-issue publication date (see <http://rnajournal.cshlp.org/site/misc/terms.xhtml>). After 12 months, it is available under a Creative Commons License (Attribution-NonCommercial 4.0 International), as described at <http://creativecommons.org/licenses/by-nc/4.0/>.

Email Alerting Service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).

20% OFF Functional LncRNA PCR Arrays
Explore LncRNA functions and their biomarker potential



To subscribe to RNA go to:
<http://rnajournal.cshlp.org/subscriptions>
